Journal of Immunological Methods, 161 (1993) 279–280

© 1993 Elsevier Science Publishers B.V. All rights reserved 0022-1759/93/$06.00

Contents to volume 161 (1993)

Number 1 – 5 May 1993

Research reports

Europium-labelled recombinant protein G. A fast and sensitive universal immunoreagent for time-resolved immunofluorometry
E. Markela, T.H. Ståhlberg and I. Hemmilä (Finland) 1

Detection of human T cell receptor cDNAs (α, β, γ and δ) by ligation of a universal adaptor to variable region
Y. Tsuruta, S. Iwagami, S. Furue, H. Teraoka, T. Yoshida, T. Sakata and R. Suzuki (Japan) 7

A quantitative ELISA for dystrophin
G.E. Morris, J.M. Ellis and Nguyen thi Man (UK) 23

Simultaneous electroelution of whole SDS-polyacrylamide gels for the direct cellular analysis of complex protein mixtures
P. Andersen and I. Heron (Denmark) 29

Detection of distinct sets of newly synthesized polypeptides in supernatants of TCR-triggered T cell clones. Implication for the search for new lymphokines
G. Trenn, J. Sykora, C. Teschendorf, R. Taffs, G. Brittinger and M.V. Sitkovsky (USA, Germany) 41

Quantitation of IgE and carcinoembryonic antigen (CEA) by optical beam deflection (OBD) measurement of dot-immunobinding assay patterns visualized by an ELISA technique
S. Matsuzawa, H. Kimura, C.-Y. Tu, T. Kitamori and T. Sawada (Japan) 59

Quantitation of IL-4 expression in small numbers of cells from mice
R.A. Barthelson (USA) 67

Characterization of human tumor cell lines transduced with the cDNA encoding either tumor necrosis factor α (TNF-α) or interleukin-2 (IL-2)
J.R. Yannelli, C. Hyatt, S. Johnson, P. Hwu and S.A. Rosenberg (USA) 77

Monoclonal antibody-specific immobilisation of erythrocyte antigens (MAIEA). A new technique to selectively determine antigenic sites on red cell membranes
A.C. Petty (UK) 91

Optimization of a time-resolved immunofluorometric assay for tumor-associated trypsin inhibitor (TATI) using the streptavidin-biotin system
S. Osman, U. Turpeinen, O. Itkonen and U.-H. Stenman (Finland) 97

Antibodies to brain integral membrane proteins in systemic lupus erythematosus
J.G. Hanly and C. Hong (Canada) 107

A flow cytometric study of the membrane potential of natural killer and K562 cells during the cytotoxic process
K. Radošević, T.C. Bakker Schut, M. Van Graft, B.G. De Grooth and J. Greve (Netherlands) 119

Short communications

Preservation of phenotype and function of positively selected virus-specific $CD8^+$ T lymphocytes following anti-Fab detachment from immunomagnetic beads
A.M. Geretti, C.A.C.M. Van Els, C.A. Van Baalen, M.C.M. Poelen and A.D.M.E. Osterhaus (Netherlands) 129

Modified procedure for labelling target cells in a europium release assay of natural killer cell activity
R. Pacifici, S. Di Carlo, A. Bacosi, I. Altieri, S. Pichini and P. Zuccaro (Italy) 135

Book review 139

Announcements 141

Notes to authors 143

Number 2 – 26 May 1993

Research reports

Production of bi-specific monoclonal antibodies in a hollow-fibre bioreactor
A. Gorter, R.J. Van de Griend, J.D.H. Van Eendenburg, W.H.B. Haasnoot and G.J. Fleuren (Netherlands) 145

Detection of inhibition of HIV-1 protease activity by an enzyme-linked immunosorbent assay (ELISA)
H.-W. Mansfeld, S. Schulz, G. Grütz, R. Von Baehr and S. Ansorge (Germany) 151

Rapid generation of monoclonal antibody-secreting hybridomas against African horse sickness virus by in vitro immunization and the fusion/cloning technique
N. Stäuber, U. Kihm and K.C. McCullough (Switzerland) 157

Phage display as a rapid gene expression system: production of bioactive cytokine-phage and generation of neutralizing monoclonal antibodies
H. Gram, U. Strittmatter, M. Lorenz, D. Glück and G. Zenke (Switzerland) 169

The use of T bag synthesis with paper discs as the solid phase in epitope mapping studies
W. Van 't Hof, M. Van den Berg and R.C. Aalberse (Netherlands) 177

Mice ascites as a source of polyclonal and monoclonal antibodies
W. Mahana and A. Paraf (France) 187

The separation and identification by monoclonal antibodies of dog IgG fractions
G. Mazza, W.P.H. Duffus, C.J. Elson, C.R. Stokes, A.D. Wilson and A.H. Whiting (UK) 193

Preparation and characterization of monoclonal antibodies against ornithine decarboxylase
R.G. Schipper, R.G.J. Rutten, M. Sauerbeck, W.J.G. Schielen, P.J.H.M. Adams, J. Kopitz, P. Bohley, G.I. Tesser and A.A.J. Verhofstad (Netherlands, Germany) 205

Single and dual labelling of cytotoxic target cells. Comparison of three radioactive tracers, [^{35}S]methionine, [^{75}Se]selenomethionine, and chromium-51
J. Heymer and W. Leibold (Germany) 217

Use of immunoplot analysis for the identification of immunodominant non-variant antigens of *Trypanosoma brucei rhodesiense*
P.J. Tidey, M. Hommel and D.H. Smith (UK) 223

An MTT-based method for the in vivo quantification of myotoxic activity of snake venoms and its neutralization by antibodies
B. Lomonte, J.M. Gutiérrez, M. Romero, J. Núñez, A. Tarkowski and L.Å. Hanson (Costa Rica, Sweden) 231

Characterization of anti-HLA class II monoclonal antibody LGII-612.14 reacting with formalin fixed tissues
M. Temponi, U. Kekish, C.V. Hamby, H. Nielsen, C.C. Marboe and S. Ferrone (USA) 239

An ELISA assay for murine interleukin-1β
R.C. Newton, R. Dowling, A.J. Daulerio and S. Culp (USA) 257

Short communication

Monoclonal antibodies against human hemopexin and their application
Z. Hrkal, H. Cajthamlová, I. Kalousek and J. Bartošová (Czech Republic) 265

Dissociation of fluorescein-labelled haptens from polyclonal antibodies. The influence of hapten molecular weight
D.L. Colbert, D.S. Smith and J. Landon (UK) 269

Letter to the editors

How to demonstrate specificity of antibody binding to synthetic peptides?
W. Van 't Hof, F.J. Van Milligen, P.C. Driedijk, M. Van den Berg and R.C. Aalberse (Netherlands) 273

Erratum

Artifacts with the thiocyanate elution method for estimating relative antibody avidity (J. Immunol. Methods 157 (1993) 269)
B.M. Gray and D.R. Shaw (USA) 277

Contents to volume 161 (1993) 279

Notes to authors 281